

RIGIDITY OF THE GROUP TOPOLOGY FOR p -ADIC SEMISIMPLE GROUPS AND AUTOMORPHISM GROUPS OF TREES

RUPERT M^cCALLUM

ABSTRACT. In [7] Kramer proves for a large class of semisimple Lie groups that they admit just one locally compact σ -compact Hausdorff group topology. We show how to adapt his argument to the group of k -rational points of any connected centreless semisimple algebraic group defined over a non-Archimedean local field k . We also show how to adapt his argument to the full automorphism group of a regular locally finite bipartite tree.

In [7] Kramer explores the question of when a semisimple Lie group G admits just one locally compact σ -compact Hausdorff group topology, or, to put it another way, when it is the case that any abstract isomorphism $\varphi : \Gamma \rightarrow G$ whose domain is a locally compact σ -compact Hausdorff topological group is necessarily a homeomorphism. He proves that this is the case for a connected centreless real semisimple Lie group for which all the simple ideals in the Lie algebra are absolutely simple. In this paper we shall generalise these arguments to other groups. First we shall discuss semisimple algebraic groups over non-Archimedean local fields, then we shall discuss the full automorphism group of a regular locally finite bipartite tree.

1. SEMISIMPLE ALGEBRAIC GROUPS OVER NON-ARCHIMEDEAN LOCAL FIELDS

The key element in Kramer's argument in [7] is Theorem 7, which gives a sufficient condition for a surjective abstract group homomorphism to be both open and continuous. We need to obtain a generalisation of this result to algebraic groups defined over non-Archimedean local fields. We shall now show how this is done. But first we shall briefly discuss the notion of a non-Archimedean local field. We base our discussion on Chapter 1 of [16].

Definition 1.1. A local field is a locally compact non-discrete Hausdorff topological field.

Definition 1.2. Given any local field k , there is a Haar measure μ (unique up to multiplication by a positive constant) on the σ -algebra of Borel subsets of k , with respect to the abelian group structure of k under addition. When S is some compact set with nonempty interior, the equation $\mu(\lambda \cdot S) = \text{mod}_k(\lambda) \cdot \mu(S)$ defines the function mod_k uniquely independently of the choice of Haar measure μ , and the function mod_k so defined is a valuation. If the valuation is Archimedean (that is, it is unbounded on the smallest subring of k containing 0 and 1), then the local field is said to be Archimedean, otherwise the local field is said to be non-Archimedean.

Theorem 1.3. *The local fields are classified as follows. Every Archimedean local field is isomorphic to either $\mathbb{R}$ or $\mathbb{C}$ with the usual topology. Every non-Archimedean local field is isomorphic either to a finite extension of the p -adic number field $\mathbb{Q}_p$, with the topology coming from the unique valuation extending the p -adic valuation on $\mathbb{Q}_p$, or to the field of Laurent*

series with coefficients in a finite field, with the topology coming from the unique valuation (up to equivalence) on such a field.

Proof. See Chapter 1 of [16]. □

Now we state Theorem 3 and 4 of [7].

Theorem 1.4 (Open Mapping Theorem). *Let $\psi : G \rightarrow H$ be a surjective continuous homomorphism between locally compact Hausdorff topological groups. If G is σ -compact, then ψ is an open map.*

Proof. See Hewitt and Ross [4], II 5. 29 or Stroppel [12], 6.19. □

Remark 1.5. In the statement of the theorem in [7] the hypothesis is given as “ H is σ -compact”. This would appear to be a typographical error.

Theorem 1.6 (Automatic Continuity). *Suppose that G is a locally compact Hausdorff topological group and that H is a σ -compact Hausdorff topological group. Assume that $\psi : G \rightarrow H$ is a group homomorphism which is also a Borel map, i.e. that the preimage of every open set $U \subseteq H$ is a Borel set. Then ψ is continuous.*

Proof. This is a special case of Hewitt and Ross [4], V.22.18; see also Kleppner [6], Thm. 1. □

Next we shall prove a version of Lemma 5 of [7] which applies to absolutely simple algebraic groups over non-Archimedean local fields. Before doing this we briefly discuss the strong topology on the set of k -rational points of an affine k -variety where k is a topological field. This is discussed in the algebraically closed case in Chapter I §10 of [8], where the main properties are left as an exercise for the reader. It is also discussed in the case $k = \mathbb{C}$ in Appendix III of [15].

Definition 1.7. Suppose that V is an affine variety over a Hausdorff topological field k , and suppose that we are given a k -embedding of V in an affine space $\mathbb{A}^n(\bar{k})$, where $\bar{k}$ is an algebraic closure of k , and the affine space is given the obvious k -structure. Then the topology on k induces a topology on the set of k -rational points $\mathbb{A}^n(k)$ of the affine space $\mathbb{A}^n(\bar{k})$, and thereby induces a topology on $V(k)$, the set of k -rational points of the variety V . This topology on $V(k)$ does not depend on the choice of embedding in an ambient affine space, and is called the strong k -topology on $V(k)$.

We also make the observation that if G is an affine algebraic group over a local field k , then $G(k)$ is σ -compact in the strong topology. This is because every local field is σ -compact, so it follows that the set of k -rational points in an affine space over a local field k is also σ -compact, and therefore every closed, and in particular every Zariski closed, subset of the affine space is σ -compact. (It is easy to see that the strong topology is finer than the Zariski topology.)

Lemma 1.8. Let G be a centreless absolutely simple algebraic group of dimension $m > 0$, defined over a non-Archimedean local field k . Suppose that V is a constructible subset of G (in the Zariski topology), of dimension greater than zero and is defined over k . Then there exist elements $g_0, \dots, g_m \in G(k)$ such that $g_0 V(k) g_1 V(k) g_2 \dots g_{m-1} V(k) g_m$ is a neighbourhood of the identity in the strong topology on $G(k)$.

Proof. Let c_0 be a point in $V(k)$ with the property that the tangent space over k to V at c_0 (in the sense of algebraic geometry; see for example [1], AG § 16) has dimension equal to that of V , and in particular is greater than zero. The existence of such a point follows from Corollary 17.2 of chapter AG of [1], together with the fact that $V(k)$ is Zariski dense in V , which follows from the implicit function theorem for local fields (see [11], Corollary 4.5). Let X be a nonzero tangent vector to V at c_0 defined over k . Replacing V by $c_0^{-1}V$, we may assume that $c_0 = e$ and that $X \in \text{Lie}_k(G) = (T_e G)_k$. The group $G(k)$ acts via Ad irreducibly on $\text{Lie}_k(G)$. Thus we can find elements $a_1, \dots, a_m \in G(k)$ such that $\text{Ad}(a_1)(X), \dots, \text{Ad}(a_m)(X)$ is a basis of $\text{Lie}_k(G)$. Now, from the implicit function theorem for local fields we know that there is a diffeomorphism from any sufficiently small open neighbourhood of the identity in $G(k)$ (in the strong topology) onto an open subset of an m -dimensional vector space over k (in the vector space topology induced from the topology on k). It then follows from the inverse function theorem for local fields (see [11], Proposition 4.3) that $h_1 V(k) h_1^{-1} h_2 V(k) h_2^{-1} \dots h_m V(k) h_m^{-1}$ is a neighbourhood of the identity in $G(k)$ (in the strong topology). This proves the result. $\square$

In what follows, the topology on $G(k)$ is assumed to be the strong topology.

Lemma 1.9. Let G be a centreless absolutely simple algebraic group of dimension $m > 0$ defined over a non-Archimedean local field k . Assume that $D \subseteq G(k)$ is a compact neighbourhood of the identity. Let $U \subseteq G(k)$ be an arbitrary neighbourhood of the identity. Then there exist elements $a_1, \dots, a_m \in G(k)$ such that $[a_1, D][a_2, D] \dots [a_m, D]$ is a neighbourhood of the identity which is contained in U .

Proof. This is proved exactly as in [7]. Let $a \in G(k) - \{e\}$. Then $\text{Ad}(a) \neq \text{id}$, so there exists $X \in \text{Lie}_k(G)$ with $\text{Ad}(a)(X) - X \neq 0$. Since $G(k)$ acts irreducibly on $\text{Lie}_k(G)$, we can find elements $a_1, \dots, a_m$ in any neighbourhood of the identity and vectors $X_1, \dots, X_m \in \text{Lie}_k(G)$ such that $\text{Ad}(a_1)(X_1) - X_1, \dots, \text{Ad}(a_m)(X_m) - X_m$ is a basis of $\text{Lie}_k(G)$. It follows readily from the inverse function theorem and implicit function theorem for local fields that $[a_1, D] \dots [a_m, D]$ is a compact neighbourhood of the identity.

Let now U be an open neighbourhood of the identity. Consider the continuous map $h : (G(k))^m \times (G(k))^m \rightarrow G(k)$, $(x_1, \dots, x_m, y_1, \dots, y_m) \mapsto [x_1, y_1][x_2, y_2] \dots [x_m, y_m]$. We have that $h(\{e\}^m \times D^m) \subseteq U$. By Wallace's Lemma, see Kelley [5], 5.12, there is an open neighbourhood V of the identity such that $h(V^m \times D^m) \subseteq U$. The claim follows if we choose $a_1, \dots, a_m \in V$. $\square$

As discussed in Chapter 5 of [14], given any Zariski-connected semisimple algebraic group H defined over a field k there exists an algebraic group $\text{Aut } H$ defined over k , such that given any field extension l of k the l -rational points of $\text{Aut } H$ are the same as the l -automorphisms of H , and the identity component (in the Zariski topology) of $\text{Aut } H$ is k -isomorphic to the adjoint group of H . Thus if H is centreless then $H(k)$ may be viewed as a subgroup of $(\text{Aut } H)(k)$. In the case where H is isotropic, the abstract automorphism group of $H(k)$ is generated by $(\text{Aut } H)(k)$ and $\text{Aut } k$, as is proved in [3].

Theorem 1.10. Let Γ be a locally compact and σ -compact Hausdorff topological group. Let H be a Zariski-connected centreless absolutely simple algebraic group defined over a non-Archimedean local field k and suppose that $H \leq G \leq \text{Aut } H$. Suppose that $\varphi : \Gamma \rightarrow G(k)$ is an abstract group isomorphism. Suppose that there is a compact subset $C \subseteq G(k)$ which

contains the k -rational part of a constructible subset $V \subseteq G$ as in Lemma 1.8, and such that $\varphi^{-1}(C)$ is σ -compact. Then φ is a homeomorphism.

Proof. By Lemma 1.8, there are elements $g_0, \dots, g_m \in G(k)$ such that $D = g_0 C g_1 C g_2 \dots g_{m-1} C g_m$ is a compact neighbourhood of the identity. We note that $\varphi^{-1}(D)$ is σ -compact. Let $U \subseteq G$ be an open neighbourhood of the identity. By Lemma 1.9 we find elements $a_1, \dots, a_m \in G(k)$ such that $E_{a_1, \dots, a_m} = [a_1, D] \dots [a_m, D] \subseteq U$ is a neighbourhood of the identity. Moreover, the set $\varphi^{-1}(E_{a_1, \dots, a_m})$ is σ -compact and in particular a Borel set in Γ . If $W \subseteq G(k)$ is an arbitrary open subset, then we find a countable collection of elements $b_j, a_{1,j}, \dots, a_{m,j} \in G(k)$ such that $W = \bigcup_{j=0}^{\infty} b_j E_{a_{1,j}, \dots, a_{m,j}}$, because $G(k)$ is second countable. Each set $\varphi^{-1}(b_j E_{a_{1,j}, \dots, a_{m,j}})$ is Borel, so $\varphi^{-1}(W)$ is a Borel set in Γ . Therefore φ is a Borel map and by Lemmas 1.4 and 1.6 is a homeomorphism. $\square$

This is the sufficient condition to which we referred earlier.

Corollary 1.11. Suppose that H is a Zariski-connected centreless absolutely simple anisotropic algebraic group over a non-Archimedean local field k , and that $H \leq G \leq \text{Aut } H$. Then $G(k)$ admits just one locally compact σ -compact Hausdorff group topology.

Proof. This is a consequence of Theorem 1.10 and the fact that if G is anisotropic then it follows that $G(k)$ is compact, which is proved in [10]. $\square$

Corollary 1.12. Suppose that H is a Zariski-connected centreless isotropic absolutely simple algebraic group over a non-Archimedean local field k which is not quasisplit, and that $H \leq G \leq \text{Aut } H$. Then $G(k)$ admits just one locally compact σ -compact Hausdorff group topology.

Proof. Let S be a maximal k -split torus. By $\mathcal{DL}(S)$ we mean the derived group of the centraliser of S , which is called the anisotropic kernel. The anisotropic kernel is a semisimple group, and its k -rational part is compact in the strong topology by [10]. Also it has dimension greater than zero, and it is not abelian, and neither is the k -rational part (since G is not quasisplit). This means that we can find an $h \in \mathcal{DL}(S)(k)$ whose conjugacy class in $\mathcal{DL}(S)(k)$ is compact and has dimension greater than zero. This set is equal to the conjugacy class of h in $\mathcal{L}(S)(k)$. The set $\mathcal{L}(S)(k)$ is closed and therefore σ -compact in any locally compact σ -compact topology on $G(k)$. The result now follows by Theorem 1.10. $\square$

As in [7], to deal with the quasisplit case we need to treat the rank-one case in slightly greater generality.

Lemma 1.13. Let G be a reductive algebraic group (not necessarily Zariski-connected) over a non-Archimedean local field k , with k -rank one. Then there exists a constructible subset of G , defined over k and of dimension greater than zero, whose k -rational part is compact in the strong topology $G(k)$ and σ -compact in any locally compact σ -compact Hausdorff topology on $G(k)$.

Proof. First consider the case where G is semisimple. The non-quasisplit case can be dealt with by the argument already given.

Since G is quasisplit, all the nodes in the Tits diagrams for G are white/encircled (see [13]), and since G has k -rank one the nodes are all in one orbit of the action of the absolute Galois

group of k . So the Tits diagram must consist of disjoint unions of finitely many copies of A_1 , or disjoint unions of finitely many copies of A_2 , with all the vertices in one distinguished orbit. This means that there exists a finite Galois extension k' of k such that the Lie algebra of G over k is the image of a Lie algebra $\mathfrak{g}$ over k' under a forgetful functor, where the Tits diagram of $\mathfrak{g}$ over k' has just one copy of A_1 or A_2 , with all nodes in one distinguished orbit. It will be seen below that our arguments will apply regardless of which representative of the isogeny class of G we choose. Thus we can assume without loss of generality that the group $G(k)$ is isomorphic (as an abstract group) either to $\mathrm{SL}(2, k')$ or ${}^f\mathrm{SU}(3, k')$ where f is a hermitian form over some quadratic extension of k' with Witt index one.

In the first case there exists a compact subgroup of $G(k)$ which is the special orthogonal group for an anisotropic quadratic form and which is equal to its own centraliser, and has positive dimension. Since this subgroup is a centraliser of a set it is closed and therefore σ -compact in any locally compact σ -compact Hausdorff group topology on $G(k)$. This proves the desired proposition in this case.

In the second case it is sufficient to prove that every three-dimensional Hermitian space of Witt index one has a two-dimensional anisotropic subspace. For then we consider the stabiliser of the two-dimensional anisotropic subspace and its orthogonal complement, which can be defined to be the centraliser of an element of $G(k)$, and find a compact conjugacy class in this stabiliser of dimension greater than zero, and then the desired proposition follows in this case as well.

So we have reduced the proposition (in the case where G is semisimple) to the lemma which follows. In the case where G is reductive, the centralisers are larger, but the appropriate subgroup or conjugacy class can still be found in both cases. $\square$

Lemma 1.14. Every three-dimensional Hermitian space of Witt index one over an arbitrary non-Archimedean local field k has a two-dimensional anisotropic subspace.

Proof. This part of the argument is due to Daniel Skodlerack. We can assume without loss

of generality that the matrix of the Hermitian form is $\begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & a \end{pmatrix}$. Let l be the quadratic extension of k over which the Hermitian form is defined and ρ the nontrivial element of $\mathrm{Gal}(l : k)$. Now let b be such that $b + \rho(b) \neq 0$. We have $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \sim \begin{pmatrix} b + \rho(b) & \rho(b) \\ b & 0 \end{pmatrix} \sim \begin{pmatrix} b + \rho(b) & 0 \\ 0 & \frac{-b \cdot \rho(b)}{b + \rho(b)} \end{pmatrix} \sim \begin{pmatrix} b + \rho(b) & 0 \\ 0 & -(b + \rho(b)) \end{pmatrix}$.

If the lower right 2-by-2 block of $\begin{pmatrix} b + \rho(b) & 0 & 0 \\ 0 & -(b + \rho(b)) & 0 \\ 0 & 0 & a \end{pmatrix}$ is not anisotropic, then the form in question is equivalent to $\begin{pmatrix} b + \rho(b) & 0 & 0 \\ 0 & b + \rho(b) & 0 \\ 0 & 0 & -(b + \rho(b)) \end{pmatrix}$. If the upper left 2-by-2 block is not anisotropic then we can multiply it by an arbitrary element of k^* , and

in particular we can choose an element of k^* which is not a norm of an element of l^* , by class field theory, (see [9], Chapter V, Theorem 1.3), and then the lower right 2-by-2 block becomes anisotropic. $\square$

Theorem 1.15. *Let H be a Zariski-connected centreless absolutely simple algebraic group over a non-Archimedean local field k which is isotropic and quasisplit. Suppose that $H \leq G \leq \text{Aut } H$. Then $G(k)$ admits just one locally compact σ -compact Hausdorff group topology.*

Proof. Let $P \subseteq G$ be a next-to-minimal parabolic, with Levi group $H \subseteq P$. The group $H(k)$ is the centraliser of a certain subgroup of $G(k)$; in particular, it is closed and therefore σ -compact in any locally compact σ -compact Hausdorff topology on $G(k)$. It is now possible to apply the argument for our result Lemma 1.13 about the rank-one case to obtain a subset C of $G(k)$ which is compact, contains the k -rational part of a variety of positive dimension as required in Lemma 1.8, and is σ -compact in any group topology of the appropriate kind, so that we can apply Theorem 1.10. $\square$

Theorem 1.16. *Let H be a product of finitely many groups H_i each of which is the rational part of a Zariski-connected centreless absolutely simple algebraic group over a non-Archimedean local field k_i . Then H admits just one locally compact σ -compact Hausdorff group topology.*

Proof. Each of the groups H_i can be defined as centralizers, which are therefore closed in any group topology. And they each admit a unique locally compact σ -compact Hausdorff group topology by Corollary 1.11, Corollary 1.12, and Theorem 1.15. $\square$

2. AUTOMORPHISM GROUPS OF TREES

Next we discuss the case of the full automorphism group of a regular locally finite bipartite tree.

Theorem 2.1. *Suppose that T is a regular bipartite tree such that there exist integers $k, l > 2$ such that all vertices of type 1 have valence k and all vertices of type 2 have valence l . Let G be the full group of automorphisms of T which preserve the bipartition. Equip G with the compact-open topology. Suppose that Γ is a locally compact σ -compact Hausdorff topological group and $\varphi : \Gamma \rightarrow G$ is an abstract isomorphism. Then φ is a homeomorphism.*

Proof. Suppose that $x \in T$, and that $y_1, y_2, \dots, y_r \in \text{Edge } T$ are distinct edges incident to x , not exhausting all of the edges adjacent to x . Let S_1 be the subtree generated by x , and all vertices $x' \neq x$ such that the geodesic from x to x' starts with one of the edges y_i , and let S_2 be the subtree generated by x , and all vertices $x' \neq x$ such that the geodesic from x to x' does not start with one of the edges y_i . Suppose that $g \in G$ has the property that the fixed vertex set of g consists precisely of S_2 . Denote by C the centraliser of g in G . C is a closed subset of the stabiliser of x (since it is contained in the setwise stabiliser of S_2) and is therefore compact. The group $\varphi^{-1}(C) \leq \Gamma$ is the centraliser in Γ of an element of Γ , and is therefore closed and therefore σ -compact. The intersection of C with the pointwise stabilizer of S_1 includes the full automorphism group of S_2 . Hence the stabiliser of x is generated by finitely many conjugates of C , and it now follows by the Baire category theorem that there must exist $g_0, g_1, \dots, g_r$ belonging to the vertex stabiliser of x , for some integer $r \geq 1$, such that $D := g_0 C g_1 C \dots C g_r$ has nonempty interior. We can then assume without loss of generality that D is a neighbourhood of the identity, and contained in a vertex stabiliser. Furthermore

D is compact, and its preimage under φ is σ -compact. The sets which are intersections of finitely many translates of D have the property that any open set is a union of such sets. Furthermore there are only countably many such sets. Thus, if U is an arbitrary open subset of G , it can be written as a countable union of intersections of finitely many translates of D . Its preimage under φ is therefore Borel. It now follows from Lemmas 1.4 and 1.6 that φ is a homeomorphism. $\square$

This result together with the following observation shows that the tree can be recovered from its automorphism group considered as an abstract group.

Theorem 2.2. *Given a regular locally finite bipartite tree, the structure of the full group of automorphisms, preserving the bipartition, with the compact-open topology, as a topological group, determines the tree up to isomorphism.*

Proof. The vertices can be identified with maximal compact subgroups of the automorphism group, and we say that two vertices G_x and G_y are adjacent if their intersection is not contained in another vertex G_z . This yields the structure of the original tree. $\square$

The above result also follows from an observation made in [2], Lemma 2.6.

REFERENCES

- [1] Armand Borel. *Linear Algebraic Groups*. Springer, 1991.
- [2] Pierre Caprace and Tom de Medts. Simple locally compact groups acting on trees and their germs of automorphisms. 2011
- [3] Armand Borel and Jacques Tits. Homomorphismes ‘abstraites’ de groupes algebriques simples. *The Annals of Mathematics*, 97(3):499–571, 1973.
- [4] E. Hewitt and K. A. Ross. *Abstract harmonic analysis. Vol. I*. Second edition, Springer, Berlin, 1979.
- [5] J. L. Kelley. *General topology*. Springer, New York, 1975.
- [6] A. Kleppner. Measurable homomorphisms of locally compact groups. *Proc. Amer. Math. Soc.*, (106):391=395, 1989.
- [7] Linus Kramer. The topology of a semisimple Lie group is essentially unique. *Advances in Mathematics*, pages 2623–2633, 2011.
- [8] David Mumford. *The red book of varieties and schemes*. Springer Lecture Notes in Mathematics 1358.
- [9] Jurgen Neukirch. *Algebraic Number Theory*. Springer, 1991.
- [10] Gopal Prasad. An elementary proof of a theorem of Bruhat-Tits-Rousseau and of a theorem of Tits. *Bull. SMF*, 110:197–202, 1982.
- [11] Peter Schneider. *p-Adic Lie Groups*. Springer, 2011.
- [12] M. Stroppel. *Locally compact groups*. European Mathematical Society (EMS), Zürich, 2006.
- [13] Jacques Tits. Classification of algebraic semisimple groups. *Algebraic Groups and Discontinuous Subgroups, Proc. Sympos. Pure Math., Boulder, CO, 1965*, pages 33–62, 1966.
- [14] Jacques Tits. *Buildings of Spherical Type and finite B-N pairs*. Springer-Verlag, 1974.
- [15] André Weil. *Foundations of Algebraic Geometry*. American Mathematical Society Colloquium Publications, Volume XXIX, 1946.
- [16] André Weil. *Basic Number Theory*. Springer-Verlag Berlin Heidelberg New York, 1967.